

1

Learning Objectives

After participating in this program you should be able to....

- Recognize the occasional medical need for directed blood donors.
- Name the different types of autologous donation.
- Describe situations where a certain type of autologous donation may be optimal.



www.aabb.org

2

2

Some basic definitions

Allogeneic- What we normally think of as a volunteer blood donation; an individual from the community donates for an unknown recipient.

Directed- An individual donates for a specific recipient. The patient will typically have a known transfusion need, such as an upcoming surgery. An allogeneic donation for a known recipient.

Autologous- An individual donates for their own upcoming surgery or for a transfusion need at a later date. For a person with a very rare blood type, the red cell can be frozen in glycerol and stored for 10 years.



www.aabb.org

3

3

More on directed donation

- Relatively uncommon, although parents sometimes request to donate for their child for an upcoming surgery.
- The blood donor must meet all criteria for allogeneic donation and have negative tests for markers of infectious disease.

Additionally:

- The donor must have an ABO/RhD type that is compatible with the patient.
- If the donor is a first degree relative, the red cell must be irradiated.



www.aabb.org

4

4

Uniform Donor History Questionnaire

Full-Length Donor History Questionnaire (DHQ)


	Yes	No
Are you		
1. Feeling healthy and well today?	<input type="checkbox"/>	<input type="checkbox"/>
2. Currently taking an antibiotic?	<input type="checkbox"/>	<input type="checkbox"/>
3. Currently taking any other medication for an infection?	<input type="checkbox"/>	<input type="checkbox"/>
4. Have you taken any medications on the Medication Deferral List in the time frames indicated? (Review the Medication Deferral List.)	<input type="checkbox"/>	<input type="checkbox"/>
5. Have you read the educational materials today?	<input type="checkbox"/>	<input type="checkbox"/>



5

Considerations with directed donation

- Typically, just the red cell unit is produced.
- Higher rate of positive markers for infectious disease.
- Donor confidentiality can be more difficult to maintain compared to a non-directed allogeneic blood donor.



www.aabb.org

6

6

More considerations

- The donation may be “crossed over” into the general community supply if the unit is not required by the patient. A collection center would need a clear policy for this.
- Any directed donation requires additional mechanisms for tracking and billing the correct amount to the recipient.
- The patient will typically be charged for the collection, whether or not the unit gets transfused. If the donor is a community donor unknown to the patient, this may not apply.



So, the tracking and billing
can get messy...

www.aabb.org



7

7


Circumstances where directed donation is useful/indicated: Scenario 1

J.H. is a 66 y.o. woman with recurrent gastrointestinal hemorrhage from stomach ulcers. She has received multiple red cell transfusions in the past, and has made several alloantibodies against red blood cells. These antibodies include: anti- c, Fya, Jka, and s. Less than 1 out of 100 people will have a red cell that is compatible with the antibodies in her circulation.

We would want to ensure that the donor’s unit is specifically tracked and allocated for the patient, rather than given to a “random” recipient.




www.aabb.org





8

8

Immunoematology- red cell antigens and antibodies



Antibody-Antigen Binding



www.aabb.org

9

9

Circumstances where directed donation is useful/indicated: Scenario 2

K.L. is a 70 y.o. man who is undergoing treatment for lung cancer. The patient's ABO blood group is AB. His red cells do not express a blood group protein known as Jk3, and (through exposure to the antigen with prior transfusion), pre-transfusion testing show that he has made anti-Jk3. The number of individuals who will be compatible with his ABO type AND lack this high-incidence antigen is low (rare in white or black individuals, 0.9% in the Polynesian population).

K.L.'s siblings may also lack Jk3, and so those who qualify for donation will be tested to determine this. If they are compatible, they can serve as the directed donor. Otherwise an unrelated blood donor whose red cells are ABO compatible and negative for the JK3 antigen will be needed for K.L.'s next transfusion.



www.aabb.org




10

10

Autologous donation

Before surgery or delivery
During surgery




www.aabb.org


11

11

Pre-surgical autologous donation



- Brief history
- Why do it?
- Donor criteria
- Why not do it?
- Actual use today
- Special situations where it may be useful.




www.aabb.org

12

12

Brief history of perioperative autologous donation (PAD)

- Autologous donation prior to surgery was promoted as a response to the AIDS epidemic, and the recognition that the virus was transmissible through transfusion, in the early 1980s.
- PAD peaked in 1993, when approximately 6% of all blood collected in the U.S. was intended for autologous use.



www.aabb.org

13

13

Advantages of autologous donation

Protection against:

- Transfusion transmissible agents
- Transfusion reactions
- Antibody formation against foreign red cells, white cells, or platelets
- Transfusion associated graft-vs-host disease (white cells in the donor unit evade destruction and destroy recipient's blood cells. Rare, can be avoided by irradiation)

HIV residual risk per donated unit: 1 in 1.5 million.



www.aabb.org

14

14

Donor Criteria

- An order from the patient's physician must be obtained.
- Minimum hemoglobin concentration of 11 g/dL or hematocrit of 33%.
- Absence of conditions presenting a risk of bacteremia.
- Note: Uniform donor questionnaire does not ask questions related to transfusion-transmissible pathogens such as HIV.



www.aabb.org

15

15

Contraindications to autologous donation

Defined by the collection center, not the FDA

In general, the following conditions will result in donor deferral:

- Unstable angina
- Recent heart attack or stroke
- History of significant cardiac disease with symptoms and no recent visit to the cardiologist
- Untreated aortic stenosis

In general, more time and thought is required by collection staff/MD for autologous donors compared to allogeneic donors.



www.aabb.org

16

16

Controversial issue- storing units with infectious agents

AABB Association Bulletin #98-5 (September 18, 1995)
The ADA, HIV, and Autologous Blood Donation

“The Supreme Court recently ruled in *Bragdon v. Abbott* that asymptomatic HIV is a disability protected by the Americans with Disabilities Act....*Bragdon* may render unlawful those policies that deny HIV-positive patients the opportunity to donate preoperative autologous units for transfusion.”

- *Side note:* Bulletins are intended to serve as guidelines, rather than Standards. But AABB members generally are expected to be aware of the recommendations in the bulletins.

For HIV and Hepatitis B, the FDA requires a written statement from the receiving transfusion service and the attending physician that HIV or HBV-positive units are acceptable before they may be shipped to the transfusing facility.



www.aabb.org

17

17

Disadvantages of autologous donation

- Data indicates that if a surgery will require less than 2 units of RBCs, autologous donation mainly serves to make the donor anemic.
- Additionally, if an autologous unit was collected, the mentality appears to be transfuse it back to the patient.



www.aabb.org



18

18

This is bad because...

Reinfusion of autologous units carries some degree of risk:

- Increased chance of bacteria compared to an allogeneic unit.
- Hemolysis of unit if not stored appropriately.
- Circulatory overload of the volume-sensitive patient.
- Possibility of a mistake and infusion of an allogeneic unit.

In one study of patients undergoing elective hysterectomy, autologous donation was an independent risk factor for transfusion. Transfusions were administered to 25 out of 140 women who had given an autologous donation, compared to 1 of 123 women who had not.

Kanter MH. *JAMA* 1996;276:798.



www.aabb.org

19

19



Autologous donation is more expensive than allogeneic donation:

- Special procedures are required to ensure that the unit is tracked and issued to the correct hospital.
- Transfusion service staff must have procedures to check for, and issue, autologous units before issuing allogeneic units.
- The unit cannot be “crossed over” into general inventory, unless the collection center allows for this (and the donor has completed all criteria for allogeneic donation).




www.aabb.org

20

20

Wastage of autologous units is typically high



Total autologous collections (red cells) at Canadian Blood Services by year
BloodBrief dated 2017-4-26



www.aabb.org

21

21

New England Journal of Medicine

N Engl J Med 1995;332:719-24.

SPECIAL ARTICLE

THE COST EFFECTIVENESS OF PREOPERATIVE AUTOLOGOUS BLOOD DONATIONS

JEFF ETCHASON, M.D., LAWRENCE PETZ, M.D., EMMETT KEELER, PH.D.,
 LONI CALHOUN, M.T.(A.S.C.P.), S.B.B., STEVEN KLEINMAN, M.D., CYNTHIA SNIDER, M.P.H.,
 ARLENE FINK, PH.D., AND ROBERT BROOK, M.D., SC.D.

A decision-analysis model was used to assess the cost-effectiveness of donating autologous blood for 4 surgical procedures done at UCLA in 1992.

Cost effectiveness was expressed in dollars per quality-adjusted year of life saved.



22

High level results

-Substituting autologous for allogeneic blood resulted in little expected health benefit (0.0002 to 0.00044 in quality-adjusted year of life saved) at considerable additional cost (\$68 to \$4,783 per unit of blood).

- The additional cost of autologous blood was primarily a function of units collected but not transfused, and of a more labor-intensive donation process.

The cost-effectiveness values ranged from \$235,000 to over \$23 million per quality-adjusted year of life saved.



www.aabb.org

23


23

Actual use today

In part because current reimbursement programs (including Medicare) either:

- Deny the medical necessity of PAD or
- Ignore the well-documented increase in cost

The use of PAD is quite limited.



www.aabb.org

24

24

Special situations where autologous donation may be helpful

- The individual is identified as having a very rare blood type, either through blood donation or prenatal/pre-transfusion testing



www.aabb.org

25

25

Example 1

J.P. is a 28 y.o. woman who visits her obstetrician at 12 weeks of pregnancy with her second child. As part of the usual prenatal testing, an antibody screen is performed. The test is positive, and shows the pattern for an antibody directed against a high incidence antigen. Numerous subsequent tests are unrevealing as to the identity of the antibody.



www.aabb.org

26

26

Medical exceptions needed for this donation

- Donation during pregnancy
- Allowance for low blood count during 3rd trimester

Ideal plan is to collect and freeze 1 unit early, then collect 1 unit that can remain in liquid state until the time of delivery.



www.aabb.org

27

27

Example 2- autologous donation

A.N. is identified through blood donation as having a very rare red cell type. It is common knowledge that only a few donors in the country will match his red cell type.

- Ideally: have the donor donate subsequent units using the questionnaire and testing for allogeneic donors. Educate the donor that it is most likely he will get these units back, but that you would like to freeze 2 as allogeneic and 2 as autologous for his potential future use.




www.aabb.org

28

28

Autologous collection during surgery

Acute Normovolemic Hemodilution (ANH) and cell salvage




www.aabb.org

29

29

Acute Normovolemic Hemodilution (ANH)

- Whole blood is collected in the operating room by the anesthesiologist.
- The units are kept within the operating room.
- The patient has blood volume maintained by crystalloid solutions.
- Most often the units are transfused back before the end of the case.
- Standards exist that dictate storage requirements in, and outside of, the operating room.




www.aabb.org


30

30

Visual of ANH



Whole blood collected using a scale after induction of anesthesia.
Crystalloid solution returned to the patient at the same time.



Keeping the units attached to the patient.



www.aabb.org

31

31

Touted benefits

- Decrease exposure to allogeneic blood, which should decrease or prevent:
 - infectious disease risk
 - allergic transfusion reactions
 - alloimmunization
 - immunomodulation



www.aabb.org

32

32

Generally not used in the following situations:

- Ischemic heart disease
- Impaired kidney function
- Hemoglobin < 11g/dL
- Inability to get laboratory values back quickly
- Inadequate vascular access



www.aabb.org

33

33

How widely used is ANH?

- Most often used in cases likely to have large blood loss.
- Depends on how committed the hospital is to “bloodless medicine.”
- Requires a dedicated anesthesia team who has experience ensuring that the patient has adequate blood volume in the setting of acute anemia.
- Not reimbursed
- Most often used in planned cardiac surgery cases.



www.aabb.org

34

34

Practical and regulatory considerations

ANH requires planning because:

- The blood bags must be acquired ahead of time; a typical blood donor bag set will lead to confusion and potential clotting if the wrong bag is chosen.
- If a refrigerator will be used to store units, it must be validated.
- Training on how to label the units must be done.
- Bloodless medicine patients require that the units remain connected to the patient during the surgery.




www.aabb.org

35

35

AABB perioperative blood collection standards



Specifies items such as:

- How units are to be labeled once collected.
- Storage conditions of the units within the operating room.
- Storage conditions if the units are taken out of the operating room following surgery.



www.aabb.org

36

36

Cell salvage

- A specialized suction system with centrifuge that allows for retrieval of lost red cells off of the surgical field.
- The shed blood is centrifuged and washed with normal saline. Plasma, cell fragments, and anticoagulant should be removed with this step.
- The system requires heparin (an anticoagulant) to keep the system from clotting.



www.aabb.org

37


37

Typically used in high-blood loss surgeries

- Cardiothoracic
- Vascular
- Orthopedic
- Trauma

In one systematic review, cell salvage during cardiac and orthopedic surgery resulted in an average savings of 0.68 unit of allogeneic red cells per patient.

Cochrane Database Systematic Rev 2010;(4):CD001888




www.aabb.org

38

38

Picture of cell salvage device



www.aabb.org

39

39

Cell salvage (Cont.)

Typically, the anesthesia technician is trained on setting up and operating the equipment.

Care must be taken not to set the vacuum pressure too high or the cells will lyse.

On a practical level, Policy decisions will need made re: how the equipment and the staff can be shared among different clinical services.



www.aabb.org

40

40

More recent use- obstetrical hemorrhage

-Fear in the past about contamination with amniotic fluid and potential to cause amniotic fluid embolus.

- Current teaching is to use 2 suction devices; one for cell salvage and one (wall suction) for amniotic fluid.
- Not generally set up unless the patient is known to be at high risk for significant post-partum hemorrhage.



www.aabb.org



41

41

Transfusion medicine MD as consultant: Bloodless medicine patient

AH is a 32 year-old woman who is pregnant with twins. Due to her religious beliefs, she does not want to receive any blood products should she experience a post-partum hemorrhage.

Care plan, made between OB, anesthesia, and transfusion medicine entails:

- Special consent detailing what products are acceptable to receive.
- Ensuring patient's iron, folate, vitamin B12 levels are replete prior to delivery.
- Being ready to perform cell salvage in OR following delivery.
- Being available at the time of delivery in case questions about pharmacologic agents are raised.
- Ensuring the patient has iron stores repleted post-delivery if there is significant blood loss.



www.aabb.org

42

42

Summary

- Autologous and directed donation are not widely used.
- These donations can be very helpful to ensuring that patients with very rare blood types have blood available when needed. Typically much planning and additional communication is required compared to an allogeneic donation for a “random” recipient.
- Other blood sparing techniques are available, with cell salvage being performed significantly more often than acute normovolemic hemodilution.
- Transfusion medicine specialists, anesthesiologists, and the clinical team work together to ensure that protocols exist and teams are well-trained.



www.aabb.org

43

43

Questions?

Contact

AABB eLearning Team

eLearning@aabb.org



www.aabb.org

Program is copyright 2020 by AABB with All Rights Reserved

44

44